

Menstrual hygiene management among reproductive-aged women with disabilities in Bangladesh

Md Nuruzzaman Khan^{1,2}  | Shimlin Jahan Khanam¹ |
Atika Rahman Chowdhury¹  | Rashed Hossain³ | Md Awal Kabir⁴ |
Md Badsha Alam^{1,5} 

¹Department of Population Science, Jatiya Kabi Kazi Nazrul Islam University, Trishal, Mymensingh, Bangladesh

²Nossal Institute for Global Health, Melbourne School of Population and Global Health, University of Melbourne, Melbourne, Australia

³Department of Sociology, Jatiya Kabi Kazi Nazrul Islam University, Trishal, Mymensingh, Bangladesh

⁴Department of Social Work, Pabna University of Science and Technology, Pabna, Bangladesh

⁵Maternal and Child Health Division, ICDDR,B, Dhaka, Bangladesh

Correspondence

Md Nuruzzaman Khan, Department of Population Science, Jatiya Kabi Kazi Nazrul Islam University, Trishal, Mymensingh, Bangladesh.

Email: sumonrupop@gmail.com

Abstract

Background: Women with disabilities in low- and middle-income countries face unique challenges in managing menstruation, affecting their health, dignity, and quality of life.

Aim: This study aimed to explore menstrual hygiene management (MHM) practices among reproductive-aged women with disabilities in Bangladesh and its impact on social participation.

Methods: We analyzed data from 51,535 women from the 2019 Multiple Indicator Cluster Survey (MICS). The outcome variables were: (i) material used to manage blood flow while menstruating (appropriate, inappropriate), (ii) availability of a private place for washing and changing menstruation rags (yes, no), and (iii) impact on attendance of social activities, school, or work during menstruation (yes, no). Disability status was considered as a major explanatory variable and categorized as no disabilities, moderate disabilities, and severe disabilities. Associations of outcome variables with explanatory variables were determined using a multilevel mixed-effects binary logistic regression model adjusted for individual-level factors, household-level factors, and community-level factors.

Results: Among the sample, 2.7% reported severe disability and 19.2% moderate disabilities, with vision-related disabilities (12.2%) being the most prevalent, followed by cognitive (9.5%) and walking disabilities (8.2%). Overall, 4% reported using inappropriate materials for menstrual flow, rising to 8.0% among those with severe disabilities and 5.8% for moderate disabilities. Women with moderate to severe disabilities had 33%–47% lower odds of using appropriate materials and 34%–44% lower odds of having a private place to change at home. Additionally, women with severe disabilities were 1.66 times more likely to report that menstruation impacted their social activities.

Conclusion: Women with disabilities in Bangladesh face significant barriers to effective MHM. Addressing these issues requires targeted interventions, including

This is an open access article under the terms of the [Creative Commons Attribution-NonCommercial](https://creativecommons.org/licenses/by-nc/4.0/) License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited and is not used for commercial purposes.

© 2024 The Author(s). *Health Science Reports* published by Wiley Periodicals LLC.

disability-inclusive menstrual health education, improved access to menstrual products and sanitation facilities, and community initiatives to reduce stigma.

KEYWORDS

adverse health outcomes, Bangladesh, menstrual health, menstrual hygiene management, women with disabilities

1 | BACKGROUND

Management of menstrual health and hygiene is essential for women's well-being and empowerment. However, it remains an unmet public health need globally, particularly in low- and middle-income countries (LMICs), where approximately 500 million women and girls lack adequate facilities for menstrual management.^{1–3} This shortfall is often compounded by a lack of access to menstrual hygiene products.¹ While one-third of women worldwide lack access to basic sanitation facilities, this figure is significantly higher in LMICs, although accurate estimates are lacking.^{4,5} Moreover, cultural taboos and social stigmas surrounding menstruation exacerbate the challenges faced by women and girls in effectively managing their menstrual hygiene.^{6–8} Consequently, inadequate menstrual hygiene management (MHM) contributes to increased health risks, including urinary tract infections, secondary infertility, anxiety, depression, and decreased self-esteem.^{9,10}

Persons with disabilities (long-term impairments that can limit their full participation in society) constitute one of the largest minority groups worldwide, with approximately 15% of the global population.^{11,12} Around 80% of them live in LMICs, highlighting the disproportionate impact on these regions.^{13,14} This demographic encompasses various impairments, including physical, sensory, intellectual, and mental health disabilities. Furthermore, there is a global trend of increasing disability prevalence with particularly more pronounced in LMICs, attributed partly to population aging, the rising incidence of chronic diseases, and existing burden of higher poverty.^{15–20} The advancement of medical technology also contributes to this trend by enabling the survival of individuals who would otherwise not have survived.²¹

Persons with disabilities in LMICs often report poorer health outcomes, particularly due to limited access to health-related information.^{18,22} Mobility restrictions and lower participation in education further contribute to these disparities.^{23,24} Additionally, they often rely on social support networks or family members for their basic needs, including healthcare.^{25–27} This dependence creates challenges in prioritizing and addressing health issues due to limited options.^{28,29} Mobility restrictions further exacerbate this situation by hindering access to freely available services, typically accessible to the general population.²³

These structural issues often result in MHM for persons with disabilities being overlooked in LMICs.³⁰ Existing studies in this setting often focus on the general population and their MHM practices, with findings revealing concerning scenarios, such as the reuse of the

Key points

What is known

- Disabilities pose an ongoing challenge in low- and middle-income countries, often leading to poor health outcomes and lower utilization of healthcare services.

What is new

- We found poor menstrual health management among reproductive-aged women with disabilities.
- Participation in social activities among women with disabilities was also found to be severely affected.

Clinical implications of the study findings

- Poor menstrual health indicates a risk of facing menstrual-related health issues, including reproductive tract infections.

same old cloth for several months.^{31–33} Commonly reported barriers to safe MHM for persons with disabilities include cultural beliefs, family environment, education level, poverty, lack of appropriate sanitation facilities, cost, and access to menstrual hygiene products.^{32,33} These factors, combined with the structural barriers mentioned above, suggest very poor MHM among persons with disabilities.^{32–35} Bangladesh exemplifies this experience, although relevant studies conducted so far are lacking, with a few studies focusing on specific disability types or small areas.^{36–40} Therefore, this study aims to explore MHM practices among reproductive-aged women with disabilities in Bangladesh.

2 | METHODS

2.1 | Study setting and sampling techniques

We extracted data from the 2019 Multiple Indicator Cluster Survey (MICS) and analyzed. This nationally representative household-based cross-sectional survey was conducted by the Bangladesh Bureau of Statistics (BBS) as part of UNICEF's initiative to collect data on the well-being of mothers and children in LMICs through standardized surveys. The survey took place from January 19, 2019 to June 1,

2019 and sampled households using a two-stage stratified cluster sampling technique, covering all 64 districts of Bangladesh. Initially, 1300 Primary Sampling Units (PSUs), areas with an approximate population of 200 to 300 households, were selected based on the 2011 Bangladesh Population and Housing Census. In the second stage of sampling, 20 households were chosen from each PSU using systematic random sampling techniques, resulting in a total sample of 64,400 households. Of these, 61,602 households were occupied, and 61,242 were successfully interviewed, yielding a household response rate of 99.4%. Among the selected households, 68,711 eligible women met the inclusion criteria: reproductive-aged (15–49 years) women who either resided in the selected households as usual residents or stayed there during the most recent night. Of these eligible women, 64,378 were successfully interviewed, resulting in a response rate of 93.7%. Detailed information about the methodology and sampling structure employed in the survey has been published in the respective survey report.⁴¹

2.2 | Analyzed sample

Data from 51,535 women from the original sample who met the inclusion criteria for this study were analyzed. The inclusion criteria were as follows: (i) reproductive-aged women (18–49 years), (ii) availability of information on women who menstruated in the last 12 months and reported related information, (iii) women who reported not being able to participate in social activities, school, or work during their last menstruation in the last 12 months, and (iv) women who reported their disability status. Importantly, we considered age 18 or older for our analyzed sample because the survey collected disability-related data only from women aged 18 years or above.

2.3 | Outcome variables

We considered three aspects of MHM and related consequences as outcome variables. These included: (i) material used to manage blood flow while menstruating (appropriate, inappropriate), (ii) availability of a private place for washing and changing menstruation rags (yes, no), and (iii) impacted in attendance of social activities, school, or work during menstruation (yes, no).

Relevant information on the use of appropriate materials during the last menstruation was collected by asking women two specific questions: “*Did you use any materials such as sanitary pads, tampons, or cloth?*” Women who responded “Yes” to this question were asked, “*What type of materials did you use in your last menstruation?*” The response options were (i) sanitary napkin, (ii) cotton pad, (iii) tissue, (iv) cloth, and other. The usage of materials during the last menstruation was categorized as appropriate material (i.e., sanitary pads and new cloths) and inappropriate material (i.e., cloth, tissue, cotton pad, and other). For the availability of a private place for washing and changing at

home during menstruation, women were asked the question, “*During your last menstrual period, were you able to wash and change in privacy while at home?*” The response options were “Yes” and “No.” Furthermore, we generated a separate variable for whether menstruation impacted in participation of social activities, including schooling and work. During the survey, women were asked, “*Due to your last menstruation, were there any social activities, school, or workdays that you did not attend?*” The response options were “Yes” and “No.” We reclassified the responses as “impacted” if women responded yes and otherwise classified as “not impacted.”

2.4 | Explanatory variables

The level of disability (no disabilities, moderate disabilities, and severe disabilities), including the types of disabilities, was the primary exposure variable for this study. These variables were assessed using a set of six questions from the Washington Group Short Set of Disability based on the World Health Organization's International Classification of Functioning, Disability, and Health.⁴² The six questions encompassed all six types for assessing disability: vision, hearing, mobility, self-care, communication, and cognition-related disability. The questions were: “(1) *Do you experience difficulty with your vision, even while using glasses?* (2) *Do you experience difficulty with your hearing, even with the assistance of a hearing aid?* (3) *Do you encounter difficulty with walking or ascending stairs?* (4) *Do you face challenges with remembering or maintaining focus?* (5) *Do you find self-care activities, such as bathing, dressing, feeding, toileting, etc., to be problematic?* and (6) *Do you struggle with communication, such as comprehending or being understood?*” The possible responses were four options for each question to respond: “(a) no difficulty, (b) some difficulty, (c) a lot of difficulty, and (d) cannot do or unable to see/hear/walk/remember/self-care/communicate at all.” We categorized the disability to analyze it in two different ways. First, we generated a variable with three mutually exclusive categories: (i) no disabilities, (ii) moderate disabilities, and (iii) severe disabilities. Women were classified as “no disabilities” if they responded “no difficulty at all” to all six questions, “moderate disabilities” if women reported ‘some difficulty’ to at least one item, and “severe disabilities” if they reported “a lot of difficulty or cannot function at all” to at least one item. Additionally, we included specific variables for each type of disability individually. For each question, we generated a binary variable—“yes” if women reported having “some difficulty and a lot or cannot do at all” and “no” if they reported having “no difficulty.” We also considered types of disabilities as explanatory variables.

2.5 | Covariates

This study incorporated several covariates to assess the association between the exposure variables and outcome variables. We selected covariates through a comprehensive process. Initially, we conducted

a comprehensive search of previous literature to identify relevant variables using several databases based on LMICs and Bangladesh.^{2,8-10,26,29,30,32,35-37,39,43} Next, we ensured their availability in the MICS data set. Then we assessed them for their statistical associations with the outcome variables. In the final analysis, variables that showed statistical significance were included and classified into three categories: individual-level, household-level, and community-level factors. Individual-level factors comprised women's age groups (18–24, 25–34, and ≥35 years), educational attainment (preprimary, primary, secondary, and higher), and mass media exposure (exposed vs. not exposed). Household-level factors included the wealth quintile (poorest, second, middle, fourth, and richest). Community-level factors included the place of residence (urban vs. rural) and administrative divisions (Barishal, Chattogram, Dhaka, Khulna, Mymensingh, Rajshahi, Rangpur, Sylhet).

2.6 | Statistical analysis

Descriptive statistics were used to summarize the characteristics of the respondents. The surveys we analyzed had a nested structure, with individuals nested within households and households nested within clusters. Previous studies have shown that using simple logistic regression models on hierarchical data produces less precise results. Therefore, we determined the associations of outcome variables with explanatory variables using a multilevel mixed-effects binary logistic regression model. We ran both unadjusted and adjusted models separately for each outcome. Unadjusted models included only outcome and exposure variables, while adjusted models accounted for individual, household, and community-level covariates. Sampling weight and complex survey design were considered in all analyses. Results were presented as unadjusted (cOR) and adjusted odds ratios (aOR) with their corresponding 95% confidence intervals (95% CI). This study was designed and reported following the Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) guidelines. All statistical analyses were performed using Stata software (version 17.0).

3 | RESULTS

3.1 | Background characteristics of the respondents

Table 1 presents the background characteristics of the respondents. The mean age of the respondents was 30.8 ± 8.5 years. Around 42% of the women had secondary-level education. Nearly 70% of the total respondents reported no exposure to mass media. Approximately, 18% of women belonged to the poorest household. Geographically, a significant portion, comprising almost 76% women resided in urban areas. Moreover, 26% of the analyzed women were from Dhaka, followed by Chattogram (19.3%) as their administrative region.

TABLE 1 Background characteristics of the respondents, MICS, 2019 (weighted $N = 51,535$).

Characteristics	N	Percent	95% CI
Individual level factors			
Ages of women, mean (SD)		30.8 (8.5)	
≤24 years	14,730	28.6	28.1–29.0
25–34 years	19,906	36.7	36.2–37.2
≥35 years	17,898	34.7	34.3–35.2
Women's education			
Preprimary	8123	15.8	15.3–16.2
Primary	12,312	23.9	23.4–24.4
Secondary	21,416	41.6	41.0–42.1
Higher	9683	18.8	18.2–19.4
Exposure to mass media			
Not exposed	15,519	30.1	29.4–30.9
Exposed	36,015	69.9	69.1–70.7
Household level factors			
Wealth index			
Poorest	9007	17.5	16.8–18.2
Second	9602	18.6	18.1–19.2
Middle	10,271	19.9	19.4–20.5
Fourth	10,947	21.2	20.6–21.9
Richest	11,709	22.7	21.9–23.5
Community level factors			
Place of residence			
Urban	12,341	24.0	23.4–24.5
Rural	39,194	76.0	75.5–76.6
Divisions			
Barishal	2754	5.3	5.2–5.5
Chattogram	9941	19.3	18.9–19.7
Dhaka	13,346	25.9	25.4–26.4
Khulna	5938	11.5	11.2–11.8
Mymensingh	3489	6.8	6.5–7.0
Rajshahi	6736	13.1	12.8–13.4
Rangpur	5652	11.0	10.7–11.3
Sylhet	3678	7.1	6.7–7.7

3.2 | Distribution of disability and its types

The percentage distribution of disability levels and types is presented in Table 2. Severe disability was reported by 2.7% of the total sample, followed by 19.2% reporting moderate disability. Among specific domains of disabilities, vision-related disabilities were found in 12.2%

TABLE 2 Prevalence of types of disability in Bangladesh, MICS, 2019.

	N	Percent	95% CI
Disabilities level			
No disabilities	40,284	78.2	77.7–78.6
Moderate disabilities	9876	19.2	18.7–19.6
Severe disabilities	1375	2.7	2.5–2.8
Types of disabilities			
Vision-related disabilities			
No	45,233	87.8	87.4–88.1
Yes	6302	12.2	11.9–12.6
Hearing-related disabilities			
No	50,286	97.6	97.4–97.7
Yes	1249	2.4	2.3–2.6
Walking disabilities			
No	47,310	91.8	91.5–92.1
Yes	4225	8.2	7.9–8.5
Cognitive disabilities			
No	46,645	90.5	90.2–90.8
Yes	4890	9.5	9.2–9.8
Self-care-related disabilities			
No	50,964	98.9	98.8–99.0
Yes	571	1.1	1.0–1.2
Communication-related disabilities			
No	51,190	99.3	99.3–99.4
Yes	344	0.7	0.6–0.8

of the total sample, followed by cognitive disabilities (9.5%) and walking disabilities (8.2%).

3.3 | Distribution of menstrual hygiene management practice and its consequences in general and among women with disabilities

We presented the percentages of MHM, both in general and among women with disabilities, in Table 3 and Figure 1. Approximately 4% of the total respondents reported inappropriate use of materials while menstruating to manage blood flow, followed by around 3% reporting having no private place for washing and changing menstrual materials. About 7.5% of the total sample reported that their social activities were impacted because of menstruation. These percentages increased for women with disabilities, with higher percentages observed among those with severe disabilities followed by moderate disabilities and no disabilities. A total of 8% of women with severe

disabilities reported inappropriate use of materials to manage blood flow, followed by 6.5% reporting having no private place to wash menstrual materials, and 11.0% reporting that their social lives were impacted because of menstruation.

3.4 | Distribution of level of disabilities across selected covariates

The distribution of disabilities level across considered covariates are presented in S-Table 1. We found the highest prevalence of moderate and severe disabilities among women aged 35 and older. Women with no formal education and not exposed to mass media exhibited a higher prevalence of moderate and severe disabilities. In terms of household-level factors, women who resided in the poorest wealth quintile reported a higher prevalence of both sorts of disabilities. Moreover, women who were from Barishal, Khulna, and Mymensingh divisions had the highest prevalence of both sorts of disabilities.

Association between MHM and its consequences with the level of disabilities and disabilities types were assessed in both unadjusted and adjusted multilevel mixed effect logistic regression models. The unadjusted model revealed decreased likelihoods of appropriate MHM practices and their consequences with increasing severity and types of disabilities (S-Table 2). These findings remained consistent after adjusting for socio-demographic factors of the respondents (Table 4 and S-Table 3). We found 33%–47% lower likelihood of using appropriate materials to manage menstrual flow among women with higher levels of disabilities compared to those without disabilities. Similarly, women with moderate to severe levels of disabilities reported 34%–54% lower likelihood of having access to private spaces at home for washing and changing menstrual rags compared to women without disabilities. Moreover, women with severe disabilities were 1.66 times more likely (95% CI, 1.35–2.03) to report that their participation in social activities, school, or work was impacted during menstruation, compared to women without disabilities. Among the adjusted factors, older age was associated with inappropriate MHM practices, lack of private spaces at home for changing menstrual rags, and reduced attendance in social activities. Additionally, women with higher education levels and those residing in wealthier households, urban areas, and the Sylhet division reported higher likelihoods of practicing positive MHM behaviors. These reported findings were found consistent once we segregate these results across types of disabilities (S-Tables 4–6).


4 | DISCUSSION

The aim of this study was to explore the effects of disabilities on MHM and its associated adverse consequences in Bangladesh. Approximately 22% of our analyzed sample had a disability, with 2.7% classified as severe disabilities and 19.2% as moderate disabilities. Notably, 3.9% of the overall sample reported using inappropriate materials to manage menstrual flow. This issue is even more prevalent among women with disabilities, reaching 8.0% for

TABLE 3 Menstrual hygiene management practice in general and across persons with disabilities.

Outcomes	Overall percentage (95% CI)	Disability status, % (95% CI)			p Value +
		Women with severe disabilities	Women with moderate disabilities	Women with no disabilities	
Material used to manage blood flow while menstruating					
Not appropriate	3.9 (3.7–4.1)	8.0 (6.5–9.7)	5.8 (5.3–6.3)	3.3 (3.0–3.5)	<0.01
Appropriate	96.1 (95.9–96.3)	92.0 (90.3–93.5)	94.2 (93.7–94.7)	96.8 (96.5–97.0)	
Availability of private place for washing and changing while at home					
No	3.2 (3.0–3.4)	6.5 (5.1–8.3)	4.3 (3.9–4.8)	2.9 (2.6–3.1)	<0.01
Yes	96.8 (96.6–97.0)	93.5 (91.8–94.9)	95.7 (95.2–96.1)	97.2 (96.9–97.4)	
Menstruation impacted in participation of social activities, including schooling and work					
Not impacted	92.5 (92.2–92.7)	89.0 (87.3–90.6)	93.1 (92.5–93.6)	92.4 (92.1–92.7)	<0.01
Impacted	7.5 (7.3–7.8)	11.0 (9.5–12.7)	6.9 (6.4–7.5)	7.6 (7.3–7.9)	

Note: +p values were obtained from chi-square test in accessing differences between menstrual hygiene management practice and consequences across level of disabilities.

**FIGURE 1** Distribution of menstrual hygiene management across level of disabilities. Association of level of disabilities with menstrual hygiene management and its consequences.

those with severe disabilities and 5.8% for those with moderate disabilities. Furthermore, 6.5% of women with severe disabilities reported not having a private place for washing and changing menstrual rags at home, while 11% reported their social activities being impacted by menstruation. After adjusting for socio-demographic factors, we found significantly lower odds (33%–47%) of not using appropriate materials to manage menstrual flow among women with

moderate to severe disabilities compared to those without disabilities. Similarly, the likelihood of having a private place to change menstrual rags at home was 34%–44% lower among women with moderate to severe disabilities. Additionally, women with severe disabilities had a 1.66 times higher likelihood of reporting that their social activities were impacted by menstruation compared to women without disabilities. These findings are robust, being derived from a

TABLE 4 Adjusted associations from multilevel mixed-effects logistic regression model of assessing associations of menstrual hygiene management and its consequences with disabilities label and type of disabilities adjusted for individual, household, and community level factors in Bangladesh, 2019 ($N = 51,535$).

Exposures	Appropriate material used to manage blood flow while menstruating aOR (95% CI)+	Availability of a private place at home for washing and changing menstruation rags (yes, no) aOR (95% CI)+	Impacted in attendance of social activities, school, or work during menstruation aOR (95% CI)+
Disability level			
No disabilities (ref)	1.00	1.00	1.00
Moderate disabilities	0.67 (0.60–0.75)***	0.66 (0.58–0.76)***	0.91 (0.82–1.01)
Severe disabilities	0.53 (0.42–0.67)***	0.46 (0.35–0.59)***	1.66 (1.35–2.03)***
Domains of disabilities			
Vision-related disabilities	0.66 (0.58–0.75)***	0.80 (0.69–0.94)***	0.83 (0.74–0.94)***
Hearing-related disabilities	0.78 (0.61–0.98)**	0.51 (0.39–0.65)***	1.36 (1.09–1.70)***
Walking disabilities	0.71 (0.61–0.82)***	0.80 (0.67–0.96)**	0.76 (0.65–0.88)***
Cognitive disabilities	0.82 (0.70–0.95)***	0.47 (0.40–0.54)***	1.26 (1.12–1.43)***
Self-care-related disabilities	0.71 (0.50–1.02)	0.53 (0.36–0.77)***	0.97 (0.68–1.38)
Communication-related disabilities	0.93 (0.55–1.56)	0.36 (0.23–0.57)***	1.20 (0.80–1.82)

Abbreviations: aOR, adjusted odds ratios; CI, confidence interval.

Note: *Models were adjusted with women's age, education, mass media exposure, wealth index, place of residence, and divisions.

*** $p < 0.01$; ** $p < 0.05$.

large-scale nationally representative survey and adjusting for a broad range of socio-demographic factors. This study underscores the significant challenges faced by women with disabilities in managing menstrual hygiene and highlights the importance of tailored interventions to address their specific needs.

The findings indicate a significant disparity in menstrual hygiene management among women with disabilities compared to those without. Specifically, women with severe disabilities were about three times more likely, and those with moderate disabilities were around twice as likely, to report not using appropriate menstrual products. These observations are consistent with available literature in LMICs and Bangladesh.^{30,43,44} This suggests multiple potential underlying issues. Accessibility and availability are critical as women with disabilities may face difficulties accessing menstrual products due to physical barriers, lack of transportation, or unavailability of products in nearby locations.^{36,37,45} Economic factors further complicate this issue, as disabilities can impact economic stability, making it harder for women to afford menstrual products.⁴⁶ Education and awareness are often lacking among women with disabilities, with societal neglect in providing inclusive health education.⁴⁷ Furthermore, societal stigma and misconceptions surrounding disabilities and menstruation may contribute to a lack of awareness and education on proper menstrual hygiene practices among this population group.^{7,48} Stigma and discrimination around menstruation and disabilities also result in inadequate support from families and communities, limiting effective menstrual health management. Additionally, healthcare systems might not be adequately equipped or sensitive to the needs of women with disabilities, leading to insufficient guidance and support.^{49,50}

Women with disabilities are also more likely to report a lack of private spaces at home where they can wash and change menstrual rags. This observation is closely reflected in Bangladeshi culture and is comparable to findings from previous studies in Bangladesh and other LMICs.^{36,37,51} In a typical Bangladeshi household, family members usually share a common toilet, which women also use for changing menstrual rags and washing them, with no separate space dedicated to these activities.⁵² While women with no disabilities can manage by washing menstrual rags in the toilet and drying them in hidden areas within the household, women with disabilities face additional challenges due to restricted mobility. This issue is particularly challenging for women with disabilities who need assistance to go to the toilet, as they lack the privacy and autonomy to manage their menstrual hygiene independently.³⁶ The situation is exacerbated by limited access to accessible sanitation facilities and the cultural stigma surrounding menstruation, which further restricts the options available to these women.^{35,37}

This study found that women with disabilities are more likely to report that their social activities are impacted by their menstrual periods compared to women without disabilities. This broader impact can be attributed to several factors. First, women with disabilities often face mobility challenges, which can make it difficult to manage menstrual hygiene effectively. Second, the lack of accessible and private facilities for changing menstrual products exacerbates the problem, leading to greater social isolation during menstruation. Additionally, societal stigma and lack of awareness about the needs of women with disabilities can contribute to their social exclusion. In many LMICs, including Bangladesh, these

issues are intensified by inadequate infrastructure and support systems.^{2,27,36,37}

The findings of this study have several policy implications. Poor MHM among women with disabilities indicates the need for targeted interventions to improve accessibility and availability of menstrual hygiene products and facilities. Policies should focus on providing accessible sanitation infrastructure, ensuring economic support for women with disabilities, and promoting inclusive health education that addresses menstrual hygiene management. Raising awareness and reducing stigma around menstruation and disabilities within communities is crucial. To address this, the government should train community family planning workers. Additionally, ongoing programs for persons with disabilities should prioritize sexual and reproductive healthcare services. Additionally, healthcare systems should be equipped to support the specific needs of women with disabilities, ensuring they receive adequate guidance and assistance for MHM.

This study possesses several strengths alongside a few limitations. One of its primary strengths is the extensive reach of the MICS, facilitating the inclusion of diverse geographic regions and socio-demographic groups, thereby enhancing the representativeness of findings. Furthermore, the cross-sectional nature of the survey enables the collection of data from a large sample within a relatively short timeframe, providing researchers with a snapshot of MHM practices and challenges across the country. The data analysis was conducted comprehensively using sophisticated statistical methods, considering a range of socio-demographic characteristics of the respondents. This comprehensive approach offers insights into the prevalence of various MHM practices among women with disabilities, illuminating both common trends and disparities across different population groups. However, the cross-sectional design of this study inherently limits the ability to establish causal relationships or track changes over time. Additionally, reliance on self-reported data may introduce bias, as participants may underreport or overreport certain behaviors or experiences related to menstrual hygiene due to social desirability or recall bias. Moreover, while the MICS aims to be nationally representative, marginalized groups or individuals with disabilities may still be underrepresented or excluded from the survey sample, potentially constraining the generalizability of findings to the entire population of women with disabilities in Bangladesh. Furthermore, despite considering various factors in the model, social norms and stigma affecting MHM were not incorporated due to a lack of relevant data in the survey. These limitations underscore the necessity for complementary qualitative research and targeted interventions to augment insights gained from the MICS data, ensuring a more nuanced understanding of the intricate interplay between disabilities and MHM practices in Bangladesh.

5 | CONCLUSION

The findings indicate that women with disabilities, particularly those with severe disabilities, are more likely to encounter obstacles in accessing appropriate menstrual products and private spaces for hygiene management. Additionally, their social activities are

disproportionately affected by menstruation, highlighting the multifaceted impact of disabilities on menstrual health. These findings underscore the urgent need for tailored interventions to address the specific needs of women with disabilities in Bangladesh. Accessibility and availability of menstrual hygiene products and facilities must be improved, alongside efforts to enhance economic support and inclusive health education. Raising awareness and reducing stigma surrounding menstruation and disabilities within communities is essential to promote social inclusion and support networks for women with disabilities.

AUTHOR CONTRIBUTIONS

Md Nuruzzaman Khan: Conceptualization; supervision; formal analysis; validation. **Shimlin Jahan Khanam:** Writing—original draft; validation; writing—review and editing. **Atika Rahman Chowdhury:** Writing—review and editing; formal analysis. **Rashed Hossain:** Writing—review and editing; validation. **Md Awal Kabir:** Writing—review and editing; validation. **Md Badsha Alam:** Conceptualization; formal analysis; software; writing—review and editing; validation.

ACKNOWLEDGMENTS

We extend our sincere thanks to UNICEF MICS for providing access to the 2019 BMICS data used for this study. Additionally, we are grateful for the support from the Department of Population Science at Jatiya Kabi Kazi Nazrul Islam University, Bangladesh, where the research was carried out. We also acknowledge ICDDR,B and its recognition of the Governments of Bangladesh and Canada for their core and unrestricted funding that supports its operations and research.

CONFLICT OF INTEREST STATEMENT

The authors declare no conflict of interest.

DATA AVAILABILITY STATEMENT

The data that support the findings of this study are available from Multiple Indicator Cluster Survey. Restrictions apply to the availability of these data, which were used under license for this study. Data are available from <https://mics.unicef.org/surveys> with the permission of Multiple Indicator Cluster Survey. The data sets used and analyzed in this study are available from the UNICEF MICS Archive website: <https://mics.unicef.org/surveys>

TRANSPARENCY STATEMENT

The lead author Md Nuruzzaman Khan affirms that this manuscript is an honest, accurate, and transparent account of the study being reported; that no important aspects of the study have been omitted; and that any discrepancies from the study as planned (and, if relevant, registered) have been explained.

ORCID

Md Nuruzzaman Khan  <http://orcid.org/0000-0002-4550-4363>

Atika Rahman Chowdhury  <http://orcid.org/0009-0001-4371-085X>

Md Badsha Alam  <http://orcid.org/0000-0002-8676-6570>

REFERENCES

- Unicef. Menstrual hygiene. <https://www.unicef.org/wash/menstrual-hygiene#:~:text=Gender%20inequality%2C%20cultural%20taboos%20and,health%20needs%20to%20go%20unmet.&text=Every%20month%2C%201.8%20billion%20people,in%20a%20dignified%2C%20healthy%20way>
- Patel K, Panda N, Sahoo KC, et al. A systematic review of menstrual hygiene management (MHM) during humanitarian crises and/or emergencies in low- and middle-income countries. *Front Public Health*. 2022;10:1018092.
- Bank W WASH (Water, Sanitation & Hygiene) and COVID-19. WASH interventions for effective COVID19 pandemic response. 2020. <https://www.worldbank.org/en/topic/water/brief/wash-water-sanitation-hygiene-and-covid-19>
- WaterAid. 1 in 3 women lack safe toilets n.d. <https://www.wateraid.org/us/media/1-in-3-women-lack-safe-toilets>
- Kaveri M. Water and sanitation crises hit women and girls harder than men 2023. <https://healthpolicy-watch.news/water-sanitation-hit-women-girls-harder>
- Malik M, Hashmi A, Hussain A, et al. Experiences, awareness, perceptions and attitudes of women and girls towards menstrual hygiene management and safe menstrual products in Pakistan. *Front Public Health*. 2023;11:1242169.
- Maulingin-Gumbaketi E, Larkins S, Whittaker M, Rembeck G, Gunnarsson R, Redman-MacLaren M. Socio-cultural implications for women's menstrual health in the pacific island countries and territories (PICTs): a scoping review. *Reprod Health*. 2022;19(1):128.
- Garg S, Anand T. Menstruation related myths in India: strategies for combating it. *J Family Med Prim Care*. 2015;4(2):184–186.
- Daniel N, Kejela G, Fantahun F, Desalegn M, Guteta F. Menstrual hygiene management practice and its associated factors among in-school adolescent girls in Western Ethiopia. *Contracept Reprod Med*. 2023;8(1):1.
- Nabwera HM, Shah V, Neville R, et al. Menstrual hygiene management practices and associated health outcomes among school-going adolescents in rural Gambia. *PLoS One*. 2021;16(2):e0247554.
- Factsheet on Persons with Disabilities | United Nations Enable. Persons with Disabilities. n.d. <https://www.un.org/development/desa/disabilities/resources/factsheet-on-persons-with-disabilities.html>
- World Health Organization. Disability. 2020. https://www.who.int/health-topics/disability#tab=tab_1
- Carew MT, Braathen SH, Swartz L, Hunt X, Rohleder P. The sexual lives of people with disabilities within low- and middle-income countries: a scoping study of studies published in English. *Glob Health Action*. 2017;10(1):1337342.
- Saran A, White H, Kuper H. PROTOCOL: effectiveness of interventions for people with disabilities in low- and middle-income countries—an evidence and gap map. *Campbell Syst Rev*. 2019;15(1-2):e1006.
- Raina P, Gilsing A, Mayhew AJ, Sohel N, van den Heuvel E, Griffith LE. Individual and population level impact of chronic conditions on functional disability in older adults. *PLoS One*. 2020;15(2):e0229160.
- Sousa RM, Ferri CP, Acosta D, et al. Contribution of chronic diseases to disability in elderly people in countries with low and middle incomes: a 10/66 dementia research group population-based survey. *Lancet*. 2009;374(9704):1821–1830.
- Martin LG, Freedman VA, Schoeni RF, Andreski PM. Trends in disability and related chronic conditions among people ages fifty to sixty-four. *Health Aff*. 2010;29(4):725–731.
- Banks LM, Kuper H, Polack S. Poverty and disability in low- and middle-income countries: a systematic review. *PLoS One*. 2017;12(12):e0189996.
- Barbareschi G, Carew MT, Johnson EA, Kopi N, Holloway C. When they see a wheelchair, they've not even seen me"—factors shaping the experience of disability stigma and discrimination in Kenya. *Int J Environ Res Public Health*. 2021;18(8):4272.
- Mascayano F, Armijo JE, Yang LH. Addressing stigma relating to mental illness in low- and middle-income countries. *Front Psychiatry*. 2015;6:38.
- Rana MS, Alam MB, Khanam SJ, Kabir MI, Khandaker G, Khan MN. Prevalence and patterns of comorbidities in people with disabilities and their associated socio-demographic factors. *Sci Rep*. 2024;14(1):1425.
- Gudlavalleti VSM. Challenges in accessing health care for people with disability in the South Asian context: a review. *Int J Environ Res Public Health*. 2018;15(11):2366.
- Tshaka B, Visagie S, Ned LY. Non-use of healthcare services among persons with mobility impairments in Cofimvaba, South Africa. *Afr J Disabil*. 2023;12:1112.
- Arul U. Access to healthcare facilities among the disabled people towards the public healthcare institution. *Int J Sci Technol Res*. 2020;9(3):4334–4338.
- Devassy SM, Scaria L, Yohannan SV, Pathrose SI. Protective role of social networks for the well-being of persons with disabilities: results from a state-wide cross-sectional survey in Kerala, India. *Int J Environ Res Public Health*. 2023;20(5):4213.
- Hashemi G, Wickenden M, Bright T, Kuper H. Barriers to accessing primary healthcare services for people with disabilities in low and middle-income countries, a meta-synthesis of qualitative studies. *Disabil Rehabil*. 2022;44(8):1207–1220.
- Hunt X, Bradshaw M, Vogel SL, et al. Community support for persons with disabilities in low- and middle-income countries: a scoping review. *Int J Environ Res Public Health*. 2022;19(14):8269.
- Clemente KAP, Silva SV, Vieira GI, et al. Barreiras ao acesso das pessoas com deficiência aos serviços de saúde: uma revisão de escopo. *Rev Saude Publica*. 2022;56:64.
- Sharma S, Mehra S. Prioritizing the needs of young people with disabilities in low- and middle-income countries: an evidence-based summary. *J Family Med Prim Care*. 2020;9(2):470–472.
- Wilbur J, Torondel B, Hameed S, Mahon T, Kuper H. Systematic review of menstrual hygiene management requirements, its barriers and strategies for disabled people. *PLoS One*. 2019;14(2):e0210974.
- Wasan Y, Baxter JAB, Rizvi A, et al. Practices and predictors of menstrual hygiene management material use among adolescent and young women in rural Pakistan: A cross-sectional assessment. *J Glob Health*. 2022;12:04059.
- Rohatgi A, Dash S. Period poverty and mental health of menstruators during COVID-19 pandemic: lessons and implications for the future. *Front Global Women's Health*. 2023;4:1128169.
- Hussein J, Gobena T, Gashaw T. The practice of menstrual hygiene management and associated factors among secondary school girls in eastern Ethiopia: the need for water, sanitation, and hygiene support. *Women's Health*. 2022;18:17455057221087871.
- BBS. National Survey on Persons with Disabilities (NSPD) 2019 Project: Bangladesh Bureau of Statistics 2020.
- World Bank Group. Menstrual Health and Hygiene. 2023. <https://www.worldbank.org/en/topic/water/brief/menstrual-health-and-hygiene>
- Afiaz A, Biswas RK. Awareness on menstrual hygiene management in Bangladesh and the possibilities of media interventions: using a nationwide cross-sectional survey. *BMJ Open*. 2021;11(4):e042134.
- Alam MU, Luby SP, Halder AK, et al. Menstrual hygiene management among Bangladeshi adolescent schoolgirls and risk factors affecting school absence: results from a cross-sectional survey. *BMJ Open*. 2017;7(7):e015508.
- Murshid ME, Shimpuku Y, Islam MZ, Rahman MM, Chen S. Effects of the involvement of male counterparts in the menstrual hygiene management of women and adolescent girls with disabilities in

- selected sub-districts of Bangladesh: protocol for a quasi-experimental study. *Cureus*. 2023;15(10):e47704.
39. Power R, Wiley K, Muhit M, et al. 'Flower of the body': menstrual experiences and needs of young adolescent women with cerebral palsy in Bangladesh, and their mothers providing menstrual support. *BMC Womens Health*. 2020;20:160.
 40. Chowdhury MA. MHM for women and girls with disabilities in Dhaka.
 41. Bangladesh Bureau of Statistics (BBS) and UNICEF Bangladesh. Progotir Pathay, Bangladesh Multiple Indicator Cluster Survey 2019. Dhaka, Bangladesh: Bangladesh Bureau of Statistics (BBS); 2019.
 42. World Health Organization. *International Classification of Functioning, Disability, and Health: Children & Youth Version: ICF-CY*. World Health Organization; 2007.
 43. WaterAid. Menstrual Health and Hygiene Management for Persons with Disabilities, 2022.
 44. Wilbur J, Poilapa R, Morrison C. Menstrual health experiences of people with intellectual disabilities and their caregivers during Vanuatu's humanitarian responses: a qualitative study. *Int J Environ Res Public Health*. 2022;19(21):14540.
 45. Fahmidul Haque AKM. Knowledge, approach and status of early marriage in Bangladesh. *Sci J Pub Health*. 2014;2(3):165.
 46. Rossouw L, Ross H. Understanding period poverty: socio-economic inequalities in menstrual hygiene management in eight low- and middle-income countries. *Int J Environ Res Public Health*. 2021;18(5):2571.
 47. The Lancet Public H. Education: a neglected social determinant of health. *Lancet Publ Health*. 2020;5(7):e361.
 48. Olson MM, Alhelou N, Kavattur PS, Rountree L, Winkler IT. The persistent power of stigma: a critical review of policy initiatives to break the menstrual silence and advance menstrual literacy. *PLOS Global Pub Health*. 2022;2(7):e0000070.
 49. Mheta D, Sibiya MN, Nkosi PB. Experiences of women with disabilities in accessing maternal healthcare services: A South African case study. *Int J Environ Res Pub Health*. 2023;20(21):6966.
 50. Matin BK, Williamson HJ, Karyani AK, Rezaei S, Soofi M, Soltani S. Barriers in access to healthcare for women with disabilities: a systematic review in qualitative studies. *BMC Womens Health*. 2021;21(1):44.
 51. WaterAid. Menstrual Health & Hygiene. 2020.
 52. Nelson KB, Karver J, Kullman C, Graham JP. User perceptions of shared sanitation among rural households in Indonesia and Bangladesh. *PLoS One*. 2014;9(8):e103886.

SUPPORTING INFORMATION

Additional supporting information can be found online in the Supporting Information section at the end of this article.

How to cite this article: Khan MN, Khanam SJ, Chowdhury AR, Hossain R, Kabir MA, Alam MB. Menstrual hygiene management among reproductive-aged women with disabilities in Bangladesh. *Health Sci Rep*. 2024;7:e70145. doi:10.1002/hsr.2.70145